

IRAM Newsletter

Number 31

April 11, 1997

Calendar

15-17 April 1997: "The Far InfraRed and Submillimetre Universe", An ESA/IRAM Symposium devoted to the Far InfraRed and Submillimetre Telescope (FIRST) cornerstone mission. IRAM will be responsible for the local organisation.

18-19 April 1997: Programme Committee Meeting

29-30 April 1997: Scientific Advisory Committee Meeting

16-17 June 1997: Executive Council Meeting

September 8th, 1997: Deadline for the submission of observing proposals for the period Nov. 15, 1997 to May 15, 1998.

Please Note

In order to reduce costs we are now sending paper copies of this Newsletter to astronomical libraries only. Individuals are invited to retrieve the IRAM Newsletter by anonymous ftp (<ftp://iram.fr/dist/newsletter>) or from our Web site (<http://iram.fr/newsletter>); please check last page for details.

Contents

Position open	1
30m Telescope	1
Interferometer	3
Software	4
Recent observations	4
Scientific results	6
New Preprints	12
Programmes Scheduled on the 30-m Telescope in 1996	13

Position open

A postdoctoral position in Astronomy is open at IRAM/Granada. *See Announcement on Page 2*

30m Telescope

NEW TELEPHONE AND FAX NUMBERS FOR PICO VELETA

The Spanish telephone company *Telefónica* changed the telephone and fax numbers for the 30m telescope in the first week of March without prior notice. The numbers for the IRAM Granada office remain unchanged. The old numbers for the telescope worked on certain days in parallel with the new numbers, but from now on, only the new numbers can be used. In addition, there will be another change of the telephone number on April 15th.

These are the new fax and telephone numbers for Pico Veleta:

(the numbers for the Granada office remain unchanged)

New fax number:

Fax +34-58-481148

Telephone number until 30 April 1997:

Tel. +34-58-481211 and +34-58-481250

Telephone number from 15 April 1997:

Tel. +34-58-482002

From April 15 to 30, we should have three telephone numbers in parallel for the 30m telescope. If you cannot reach Pico Veleta by telephone or fax, please call the Granada office to inquire (*Telefónica* may have changed the numbers again ...).

ONE WEEK OF 350 GHz OBSERVATIONS AT THE 30M TELESCOPE

The IRAM 350 GHz SIS receiver has been installed and used during the week from 4 to 11 of March, with excellent results. The receiver performance was very good, with

IRAM — Institut de Radioastronomie Millimétrique

Applications are invited for a

POSTDOCTORAL POSITION IN ASTRONOMY

at IRAM Granada/Spain, starting as soon as possible, and not later than June/July 1997.

IRAM Granada operates the 30m millimeter telescope, located 50 km from Granada in the Sierra Nevada at an altitude of 3000 m, and offices (laboratories/library/administration) in the town of Granada. The total size of the group is approximately 30 persons.

The tasks of the post-doc astronomer include:

- to participate in projects to improve the telescope system in such areas as software, receivers, backends and optics, as well as to participate in the ongoing development and improvement of observing techniques (such as spectral line on-the-fly observing and frequency switching)
- to carry out part of the regular test observations (pointing, calibration, etc.) and participate in their analysis
- regular participation in the astronomer-on-duty service at the 30 m telescope to support guest observers.

These activities will occupy at least 50% of the post-doc's time. In addition he/she has the possibility to pursue his/her own scientific projects, also in collaboration with IRAM astronomers and/or outside groups.

The applicant should have a Ph.D. in astronomy or physics and preferably some experience in observational astronomy. Knowledge in some areas related to the hardware or software of a millimeter telescope is an advantage. The ability to work at high altitude (3000 m) is essential. Good knowledge of English is required.

The initial appointment will be for a period of two years with the possibility of renewal for a third year. Applications should be submitted before April 30th, 1997, to:

Prof. Dr. M. Grewing
Director
IRAM
300 rue de la Piscine
Domaine Universitaire
F-38406 St. Martin d'Hères Cedex, France
e-mail: grewing@iram.fr

Further information can be obtained from Dr. W. Wild, Station Manager, e-mail: wild@iram.es

receiver noise temperatures of 70 K to 100 K on the sky (including optics) and a sideband rejection of about 6 dB. The weather was excellent during a large fraction of the time. System noise temperatures of 400 K to 600 K at 350 GHz were not unusual, even during daytime ! These low temperatures correspond to about 0.5 mm of precipitable water vapour, clearly demonstrating the potential of the 30m telescope site for observations at 0.8 mm wavelength.

INTERRUPTION OF THE INTERNET LINK FOR ONE WEEK

IRAM Granada is connected to the Internet via a digital (ISDN) telephone line to the University of Granada. This telephone line, operated by the Spanish telephone company *Telefónica*, was interrupted for approximately one week (March 6 to 12) due to a major failure in one of *Telefónica's* telephone exchanges. Unfortunately, *Telefónica* was not able to fix the failure quicker. In the future we will have a private link to the University of Granada and will then use the telephone link just as a backup. With this, we hope to be more independent and to be able to provide a more reliable Internet connection.

This incident – which caused some inconvenience for the visiting astronomers – emphasizes what has been mentioned here earlier:

Do not rely on the Internet for your observations! Make sure to bring all necessary files (source lists, frequencies, previous data etc.) with you, or transfer them to Granada well before your observing run.

NEW CHART RECORDER SOFTWARE

A new software for the bolometer chart recorder is available now. It does not require special hardware as the previous version, and is more reliable and flexible. The new version displays its output (> 2 channels) on a window under X-Windows and, if required, two channels on the chart recorder. A hardcopy of the display window is possible as well. The current version works for bolometer observations, and we plan to offer it also for observations with heterodyne receivers.

Wolfgang WILD

Interferometer

OBSERVATIONS

Thanks to the very good weather conditions in the last 2 months, we have been able to recover most of the delay accumulated during the fall. Almost all “A” proposals are now complete, with few exceptions. Most “B” proposals are nevertheless unlikely to be started, because of unfavourable combinations of frequency and LST range: 1.3 mm observations are difficult in daytime at this epoch of the year.

A “blind” adjustment of the panels from the inner ring of antenna 5 had been done late November. This adjustment could only be checked when the interferometer was back to the compact D configuration late March. The holography of antenna 5 performed late March yields a surface accuracy of 50 μm . 3 out of 5 antennas now have a gain of 27 Jy/K at 245 GHz.

DATA REDUCTION

Visitors coming for data reduction in Grenoble will find a number of hardware and software changes.

Concerning the hardware, 16 GBytes of disk space have been recently added to the workstation dedicated to Plateau de Bure data reduction. A second, more powerful, HP-J280 workstation with 24 GBytes of storage will be added soon. Real-time archiving is now done on CD-ROMs rather than DATs, significantly reducing the time required to load the original data set.

To fully benefit from the CD-ROM archiving, the CLIC program has been modified to handle the data and the calibration parameters in two separate files. The data file is never modified, and can reside in a read-only file system like a CD-ROM. Operation is transparent for the user. The original DAT archive is progressively converted to CD-ROMs.

A new calibration procedure has been written. The new procedure utilizes the SIC Window interface, and greatly simplifies and speeds up data reduction. “Automatic” calibration is even possible on good quality data. The procedure handles dual-frequency projects by phase-referencing the 1.3 mm data to the 3 mm ones.

Based on this procedure, and capitalizing on the experience gained with the interferometer, we are studying a method to pre-calibrate the data at the Plateau de Bure.

The “Calibration CookBook” is under revision to include a more comprehensive description of the new calibration facilities.

Stéphane GUILLOTEAU

Software

A number of small installation problems and/or bugs which appeared in the DEC96 release have been fixed in the APR97 release. The APR97 release also incorporates two internal changes: the merging of two libraries (lib-gag.a and library.a), and the “disparition” of the “old” CLASS program. The new CLASS program incorporates full support for the On-The-Fly mapping.

The APR97 release is complete for HPUX, AIX and Dec-OSF1 systems. The SIC Window facilities are still not available for SunOS, Solaris, Ultrix and Linux.

Finally, since PCs are now extremely widespread and powerful enough to handle the type of data processing required by the IRAM instruments, we have undertaken a porting of the GILDAS software to the Windows-95 operating system.

Stéphane GUILLOTEAU

Recent observations

MOLECULAR OBSERVATIONS OF COMET HALE-BOPP (C/1995 O1) WITH THE IRAM PLATEAU DE BURE INTERFEROMETER.

Report by: D. Bockelée-Morvan, N. Biver, P. Colom, J. Crovisier, E. Gérard, H. Rauer (Observatoire de Paris-Meudon), D. Despois (Observatoire de Bordeaux), J. Wink (IRAM, Grenoble), R. Moreno, G. Paubert (IRAM, Grenade), J.K Davies, W.R.F. Dent (JAC, Hilo)

The gaseous coma of Hale-Bopp was observed at the IRAM Plateau de Bure Interferometer in October and November 1996 and from March 10 to March 20 1997. The spatial resolution was 1.5'' at 1mm and 3.5'' at 3 mm, corresponding to about 1500 and 3500 km on the comet in March 1997. Emission lines of HCN, HNC, CO, H₂CO, CH₃OH, CS and H₂S were successfully detected in cross-correlation mode. High quality maps were obtained for these species. SO was not detected in interferometry. In addition SO₂ (Wink et al. Circ. 6591, 1997) and HCOOH (Wink et al. Circ. 6599, 1997) were successfully detected in auto-correlation mode, providing the first identification of these species in cometary atmospheres.

These interferometric observations have several goals. First, the spatial distribution of CO (fig. 1, H₂CO and HNC should put strong constraints on their origin. *In situ* observations aboard the Giotto spacecraft have shown that a large fraction of CO and H₂CO is released in the coma rather than from the sublimation of the nucleus.

This distributed source could be organic grains or complex species, as polymers. The observations at Plateau de Bure should be able to distinguish the fraction of CO and H₂CO coming from the nucleus and constrain the size of the distributed sources. The abundance of the very unstable species HNC in cometary nuclei is another important issue, since it should be an important diagnostic on the conditions of formation of cometary material. The non-detection of SO in interferometric mode brought the proof that SO is mainly a daughter species, i.e. a photodissociation product (Wink et al. IAU Circ. 6591, 1997). Detailed modelling is needed to establish whether SO₂ (Wink et al. IAU Circ. 6591, 1997) is its main parent. The observations will also provide a 3-D reconstruction of the morphology of the gaseous coma, the high spectral resolution giving the informations along the line of sight. Several lines of CH₃OH, CO and CS were observed to study the evolution of the excitation temperature in the coma.

The abundance of SO₂ and HCOOH in the nucleus of comet Hale-Bopp will allow to better understand the links between cometary and interstellar material and the origin of comets.

See also the Comet page on IRAM's Web page at <http://iram.fr/hale-bopp/comet.html>.

Editor's Note: The continuum emission from the nucleus has also been detected with the Plateau de Bure Interferometer by Altenhof, Wink et al. (IAU telegram 6587), allowing an estimate of the size of the nucleus (diameter \sim 45 km).




Figure 1: Map of the J(2-1) transition of CO at 230 GHz obtained on March 11, 1997. The spatial resolution is 1.5".

Scientific results

MOLECULAR GAS IN THE BARRED SPIRAL M 100. I. THE IRAM 30M MAP

M.J Sempere and S. García-Burillo

Observatorio Astronómico Nacional. Campus Universitario. Apartado 1143, E-28800 Alcalá de Henares (Madrid), Spain.

Abstract. We present a study on the distribution, kinematics and evolution of the molecular gas disk in the barred SBbc spiral M100(NGC4321). The $J=2-1$ and $J=1-0$ lines of ^{12}CO have been mapped with the IRAM 30m telescope, with resolutions $\text{HPBW}=12''$ and $\text{HPBW}=21''$, within the inner 8 kpc of the disk ($1'' = 82\text{pc}$, assuming $D_{\text{M100}} = 17.1\text{Mpc}$). Complementary $^{13}\text{CO}(1-0)$ observations of the nucleus, spiral and interarm regions are used to study the nature of H_2 gas.

CO emission traces a markedly asymmetric two spiral-arm structure stretching out from a molecular gas bar. The CO bar has a diameter $D \sim 90''$ and it is aligned with the stellar bar seen in I and K band images ($\text{PA}=110^\circ$). Arm I (II) springs off the western (eastern) side of the bar at $r \sim 50''$. Arm II is split up in two armlets (at $r \sim 40-50''$), it disappears and finally shows up at a position angle close to the major axis northern crossing ($\text{PA}=330^\circ$). The splitting starts near the inner 3:1 resonance identified by Elmegreen et al (1992) in the enhanced optical images of M100. Molecular gas in the bar is strongly concentrated in a nuclear disk of $r_{\text{nuc}} \sim 30''$ and $M(\text{H}_2)_{\text{nuc}} = 2.8 \cdot 10^9 M_\odot$.

Arms I-II display different arm-interarm contrasts (on average, $R_{a/ia}(\text{CO})$ increases with radius from 2.5 ($r \sim 30''$) up to 6.5 ($r \sim 110''$), being higher for arm I than for arm II. Also arms I-II relate differently to other spiral arm tracers, underlying the asymmetry in the disk. Comparison between CO, H I and H_α maps show that there is no *coherent* sequence in the relative location of the star formation tracers along the spiral arms. Evidences of triggering of star formation along spiral arms are poor: $R_{a/ia}(\text{H}_\alpha)$ is only $\sim 2 R_{a/ia}(\text{CO})$ and systematic offsets between H_α and CO ridges (expected to lie downstream and upstream the spiral potential minimum, respectively, assuming trailing arms inside corotation) are hard to find and at places they are absent or even inverted.

CO reveals as the best tracer of gas kinematics in the inner disk. The CO rotation curve (V_{rot}) is steeper than the curves derived from the H I and H_α data. V_{rot} reaches 200kms^{-1} in less than 1 kpc. The signature of the $m = 3$ instability has been also identified in the CO derived velocity field. The magnitude and the sign of streaming motions, associated with the spiral arms and the bar, are consistent with the CO disk to be inside corotation ($r_{\text{cor}} = 110''$). A secondary wave compression develops in the eastern side where arm II is split.

Massive star formation (MSF) is inhibited along the

gaseous bar, indicating that M100 is an *evolved* barred system (Friedli and Benz, 1995). Star formation rate (measured as the ratio $\text{SFR} = F(\text{H}_\alpha)/I_{\text{CO}}$) is lower for the nuclear disk than for the disk itself. However we suspect the measurement of SFR to be subjected to major uncertainties: the $X = N(\text{H}_2)/I_{\text{CO}}$ conversion factor might be 3 times lower than implicitly assumed and to vary within the disk (lower for the interarm than for the arms and nuclear region). Moreover extinction affects $F(\text{H}_\alpha)$ mostly in the nucleus where the classical Schmidt law breaks down paradoxically ($\text{SFR}_{\text{ND}} \sim N_{\text{gas}}^\alpha$, $\alpha \sim 1$). MSF is set on at a distance $r_c \sim 12$ kpc where $N(\text{H}_2)$ approaches the Toomre (1964) gravitational instability threshold ($N_c \sim 7M_\odot$). Also for $r < r_c$, the neutral gas is mostly in the H_2 phase. H I is underabundant in the inner 6kpc and cannot be accounted for by photodissociation of H_2 by H II regions.

Asymmetry in the observed molecular gas distribution and kinematics of M100 seems related with the three-arm structure studied by Elmegreen and collaborators. Although the $m = 2$ spiral mode is still predominant in M100, other secondary modes seem at interplay reflecting the secular evolution of the disk. Compared to M51, M100 appears as an *evolved* barred spiral.

THE EXTENDED 1.3 MM CONTINUUM EMISSION AROUND CW LEO

M. A. T. Groenewegen⁽¹⁾, W. E. C. J. van der Veen⁽²⁾, B. Lefloch⁽³⁾, A. Omont⁽⁴⁾

⁽¹⁾Max-Planck Institut für Astrophysik, Karl-Schwarzschild Straße 1, D-85740 Garching, Germany

⁽²⁾Department of Astronomy, Columbia University, 538 West 120th Street, New York, NY 10027, U.S.A.

⁽³⁾IRAM, Avda. Divina Pastora 7, Nucleo Central, E-18012 Granada, Spain

⁽⁴⁾Institut d'Astrophysique de Paris, CNRS, 98bis Boulevard Arago, F-75014 Paris, France

Abstract. We have obtained a $150 \times 150''$ map of the 1.3 mm emission around the carbon star CW Leo. Emission is detected at the 3σ level out to a distance of $50''$ from the central star. From model calculations we find that the dust density distribution is described by an approximate r^{-2} law superposed on which are possible density enhancements at about 5 and $20''$ distance from the star. This could imply phases of enhanced mass loss lasting up to several hundred years duration over the last few 1000 years. The major uncertainty in a quantitative analysis is in the contribution of molecular lines. The density enhancement at $5''$ is consistent with a recent phase of increased mass loss as inferred from CO data.

Accepted by *A&A Letters*

For preprints, contact groen@mpa-garching.mpg.de, or <http://www.mpa-garching.mpg.de/~groen/groen.html>




Figure 3: The 1.3 mm continuum map of CW Leo. The noise level is 5 mJy/beam, and contours are drawn at the 3, 4, 5 and 6 σ level. The map size is $150 \times 150''$, with north up and east to the left. Scale is in arcseconds.




Figure 2: Overlaid on gray scale images of I₁₀ and I₂₁. We plot the first-moment isovelocity contours derived from the 1–0 and 2–1 transitions of ¹²CO (figures (a)(top) and (b)(bottom), respectively). Isovelocities go from 1450 km s⁻¹ to 1750 km s⁻¹ by steps of 25 km s⁻¹. Thick contours stand for the systemic velocity $v_{\text{sys}} = 1575 \text{ km s}^{-1}$. A close view shows the different strength and degree of symmetry of the streaming motions in the southern (I) and northern (II) arms. Along arm I, streaming motions can be followed more continuously and their amplitude is stronger than in arm II. The velocity field along arm II reflects again the asymmetry of the gaseous spiral response in M100: a secondary wave compression appears in the eastern side, coinciding with the splitting of arm II. This kinematical feature, present in the 2–1 and the 1–0 maps, assesses our view that we are detecting the signature of the $m = 3$ mode.

THE CHEMICAL EVOLUTION OF PLANETARY NEBULAE

R. Bachiller⁽¹⁾, T. Forveille⁽²⁾, P. Huggins⁽³⁾, & P. Cox^(4,5)

⁽¹⁾ Observatorio Astronómico Nacional (IGN), Apartado 1143, E-28800 Alcalá de Henares, Spain

⁽²⁾ Observatoire de Grenoble, B. P. 53X, F-38041 Grenoble Cedex, France

⁽³⁾ Physics Department, New York University, 4 Washington Place, New York NY 10003, USA

⁽⁴⁾ Institut d'Astrophysique Spatiale, Bât. 121, Université de Paris XI, F-91405 Orsay Cedex, France

⁽⁵⁾ Institut d'Astrophysique de Paris, 92b bd. Arago, F-75014 Paris, France

Abstract: We report millimeter line observations of CO, ¹³CO, SiO, SiC₂, CN, HCN, HNC, HCO⁺, CS, and HC₃N to study the chemistry in planetary nebulae (PNe) with massive envelopes of molecular gas. The sample observed consists of representative objects at different stages of development in order to investigate evolutionary effects: the proto-PNe CRL 2688 and CRL 618, the young PN NGC 7027, and the evolved PNe NGC 6720 (the Ring), M4-9, NGC 6781, and NGC 7293 (the Helix).

The observations confirm that the chemical composition of the molecular gas in PNe is radically different from that in interstellar clouds and the circumstellar envelopes of Asymptotic Giant Branch (AGB) stars. There are also clear trends in the chemical evolution of the envelopes. As a star evolves beyond the AGB, through the proto-PN and PN phases, the abundances of SiO, SiC₂, CS, and HC₃N decrease, and they are not detected in the PNe, while the abundances of CN, HNC, and HCO⁺ increase dramatically. Once a PN has formed, the observed abundances in the molecular clumps of the envelope remain relatively constant, although HNC is anomalously underabundant in NGC 7027. In the evolved PNe, CN is about an order of magnitude more abundant than HCN, HNC, and HCO⁺, and the average abundance ratios are CN/HCN = 9, HNC/HCN = 0.5, and HCO⁺/HCN = 0.5. These ratios are, respectively, one, two, and three orders of magnitude higher than in the prototypical AGB envelope IRC+10216. The ¹²C/¹³C ratios are $\approx 10 - 25$, within the large range found in AGB envelopes. The chemical evolution of the envelopes likely occurs through the development of photon-dominated regions produced by the ultraviolet radiation field of the central star.

The observations also provide important information on the physical conditions in the molecular gas. Multi-line observations of CN, CO, and HCO⁺ show that the clumps which form the envelopes of the evolved PNe maintain remarkably high gas densities ($\sim \text{few} \times 10^5 \text{ cm}^{-3}$) and low temperatures ($\sim 25 \text{ K}$). These values are consistent with the idea that the clumps are in rough pressure equilibrium with the more diffuse, ionized gas and can last for a significant part of the nebular lifetime, providing the environment needed for the survival of the molecules. Thus the clumping of the gas in these PNe is an essential aspect

of both their physical and chemical evolution.

Accepted by Astron. Astrophys.

Preprints can be obtained by contacting bachiller@oan.es

THE GRAVITATIONAL LENSING NATURE OF THE CLOVERLEAF UNVEILED IN CO (7-6) LINE EMISSION

D.Alloin⁽¹⁾, S.Guilloteau⁽²⁾, R.Barvainis⁽³⁾,
R.Antonucci⁽⁴⁾, L.Tacconi⁽⁵⁾

⁽¹⁾Service d'Astrophysique, CE Saclay, 91191 Gif-sur-Yvette Cedex, France

⁽²⁾Institut de Radio Astronomie Millimétrique, 300 Rue de la Piscine, F-38406 Saint Martin d'Hères, France

⁽³⁾MIT Haystack Observatory, Westford, MA 01886, USA

⁽⁴⁾UC Santa Barbara, Physics Department, Santa Barbara, CA 93106 USA

⁽⁵⁾Max Planck Institut für extraterrestrische Physik, D-85748 Garching, Germany

Abstract. We report 0.5'' resolution imaging of the Cloverleaf quasar in the CO(7-6) line and 1.3 mm continuum, as well as 1'' resolution imaging in the CO(3-2) line, performed with the IRAM interferometer. The CO(7-6) image is clearly resolved into 4 spots, demonstrating that the CO emission is also gravitationally lensed. We do not find any convincing evidence for a velocity gradient in the maps. As compared to the optical spots from a V band HST image, some of the CO spots are found to be stretched along the Einstein ring. Using the available lens model for the Cloverleaf, this suggests an intrinsic radius of ~ 600 pc for the CO emitting region. No continuum emission is detected: the measured flux density is 5 ± 3 mJy, consistent with a spectral index larger than 3.

STRUCTURE AND KINEMATICS OF A PROTOSTAR: MM-INTERFEROMETRY OF L 1157

F.Gueth⁽¹⁾, S.Guilloteau⁽¹⁾, A.Dutrey⁽¹⁾, R.Bachiller⁽²⁾

⁽¹⁾ IRAM, 300 rue de la piscine, F-38406 St Martin d'Hères, France

⁽²⁾ OAN, Apartado 1143, E-28800 Alcalá de Henares, Spain

Abstract. We present high angular resolution (2.5'') interferometric images of ^{13}CO ($J = 1 \rightarrow 0$), C^{18}O ($J = 1 \rightarrow 0$), and $\lambda 2.7$ mm continuum emission around L1157-mm, a suspected protostar powering an energetic molecular outflow. The continuum emission consists in two distinct components. The Class 0 object L1157-mm is seen as a compact source of size $\leq 1''$ (440 AU) and mass $\simeq 0.2M_{\odot}$, which is marginally resolved and elongated perpendicular to the outflow axis. In addition, spatially-extended low-level emission is observed and likely arises at the heated edges of the cavity excavated by the outflow in the surrounding envelope. Simple modelisation shows

that such an interacting structure can explain the observed morphology and the spectral energy distribution of the source.

The line observations indicate that ^{13}CO mostly originates from the envelope and the limb-brightened edges of the outflow, whereas the C^{18}O emission is more directly associated with the compact continuum source. Weak evidence for rotation has been found. Comparison of the two line tracers shows prominent redshifted self-absorption in the ^{13}CO spectrum which is very suggestive of infall motions. The infalling medium seems to be confined in a large (a few thousands AU) flattened structure.

These observations provide a quite detailed description of the structure of a Class 0 protostar, which appears to have a complex vicinity where the outflowing and accreting phenomena are closely linked together (Fig. 4)




Figure 4: 2.5'' observations of the protostar L 1157-mm: (*Left*) $C^{18}O$ $J = 1 - 0$ (thick histogram), ^{13}CO $J = 1 - 0$ (filled) and ^{13}CO $J = 1 - 0$ interferometric + single-dish (thin) spectra observed towards the central position. (*Right*) $\lambda 2.7$ mm continuum emission (greyscale) surimposed with the integrated CO emission: the extended component of the continuum emission is associated with the edges of the molecular outflow. The white ellipse is the formal result of a fit of the compact central source.

THE INTERSTELLAR MEDIUM IN THE EDGE-ON GALAXY NGC 5907: COLD DUST AND MOLECULAR LINE EMISSION

M. Dumke⁽¹⁾, J. Braine^(2,3), M. Krause⁽¹⁾, R. Zylka⁽¹⁾, R. Wielebinski⁽¹⁾, and M. Guélin⁽³⁾

⁽¹⁾ Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany

⁽²⁾ Observatoire de Bordeaux, URA 352, CNRS/INSU, B.P. 89, F-33270 Floirac, France

⁽³⁾ Institut de Radio Astronomie Millimétrique, 300 Rue de la Piscine, F-38406 Saint Martin d'Hères, France

Abstract. We present new observations of the interstellar medium in the non-interacting edge-on galaxy NGC 5907. We have observed the $J = 2 - 1$ and $J = 1 - 0$ lines of the ^{12}CO molecule and radio continuum emission at $\lambda 1.2$ mm.

The distribution of the molecular gas (as traced by CO) shows a maximum in the central region and a ring or spiral arm at $r \sim 7$ kpc. Further analysis of the major axis distribution reveals evidence for an inner ring-like structure at $r \sim 3.5$ kpc. The kinematics can be described by rigid rotation in the inner part, a turnover at ~ 3 kpc, and differential rotation with a velocity of 230 km/s in the outer disk.

The observed continuum emission is mainly due to thermal radiation of cold dust with an average temperature of $T_d = 18$ K, with a small gradient from 20 K to 16 K from the center to the outer disk. This cold dust component is necessary to explain our results.

The dust emission closely follows the molecular gas in the central region, but is also detected at large radii where no CO can be seen (Fig. 5). In these regions the dust absorption cross section per H atom at $\lambda 1.2$ mm is estimated to be $\sigma_{\lambda}^{HI} \sim 4.5 \cdot 10^{-27}$ cm², a value similar to that in the outer parts of other galaxies.

From the $\lambda 1.2$ mm emission we estimated a molecular mass of NGC 5907 of $0.9 \cdot 10^9 M_{\odot}$, about 50 % smaller than from the CO emission. By combining the CO and continuum data we found that the CO-H₂-conversion ratio increases with galactocentric radius, from $\sim 0.7 \cdot 10^{20}$ at the centre to $\sim 1.6 \cdot 10^{20}$ cm⁻²/K km s⁻¹ at $r = 7.5$ kpc.

Accepted for publication in A&A




Figure 5: Contour map of the continuum emission of NGC 5907 at 245 GHz, overlaid onto an optical image extracted from the Digitized Sky Survey. The beam size of $15''$ is indicated by the filled white circle in the lower right corner. The rms noise depends on the location in the map; on the galaxy it is about 1.5 mJy/beam area, contour levels are 4, 8, ..., 24 mJy/beam area

EVOLUTION OF THE OUTGASSING OF COMET HALE-BOPP (C/1995 O1) FROM RADIO OBSERVATIONS.

Nicolas Biver, Dominique Bockelée-Morvan, Pierre Colom, Jacques Crovisier, John K. Davies, William R.F. Dent, Didier Despois, Eric Gérard, Emmanuel Lellouch, Heike Rauer, Raphael Moreno, Gabriel Paubert

Abstract: Since soon after its discovery at 7 AU from the Sun, comet Hale-Bopp has been studied by radio spectroscopy to investigate the gas species associated with its activity. We present an extensive monitoring performed with the Nançay, IRAM-30m and JCMT radio telescopes, showing the progressive release of CO, CH₃OH, HCN, OH, H₂S, CS, H₂CO, CH₃CN and HNC as the comet approached the Sun from 6.9 to 1.4 AU. These data provide clues to the chemical and physical state of cometary ices and allow us to investigate the fractionation processes

that accompany their sublimation upon solar heating. Simultaneous observations of the relative intensities and line shapes of several transitions are used to measure the expansion velocity and temperature of the cometary atmosphere and their evolution with heliocentric distance, as well as to study possible sublimation from icy grains. The more volatile species are relatively more abundant in the coma far from the Sun, but there is no direct correlation between overabundance and volatility. Evidences for H₂O sublimation from icy grains are seen beyond 3.5 AU from the Sun. The change from a CO-driven coma to a H₂O-driven coma occurred at about 3 AU. The gas outflow velocity and temperature increased with decreasing heliocentric distance.

Published in *Science*, **275**, 1915 (1997).

MILLIMETER INTERFEROMETRY TOWARDS THE ULTRA-COMPACT HII REGION W3(OH)

F. Wyrowski⁽¹⁾, P. Hofner⁽²⁾, P. Schilke⁽¹⁾, C.M. Walmsley^(1,3), D.J. Wilner⁽⁴⁾, J.E. Wink⁽⁵⁾

⁽¹⁾I. Physikalisches Institut, Univ. zu Köln, Zùlpicherstr. 77, D-50937 Köln, Germany

⁽²⁾Cornell University, NAIC, Arecibo Observatory, P.O. Box 995, Arecibo, P.R. 00613

⁽³⁾Osservatorio Astrofisico di Arcetri, Largo E.Fermi 5, I-50125 Firenze, Italy

⁽⁴⁾Harvard Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138, USA

⁽⁵⁾Institut de Radioastronomie Millimétrique, Domaine Universitaire, F-38406 St-Martin-d'Hères, France

Abstract: We used the IRAM Interferometer to map the J=1-0 and J=2-1 lines of C¹⁷O as well as the continuum at 112 and 225 GHz of the W3(OH) region. Towards the cluster of water masers W3(H₂O) we observed compact continuum emission at both frequencies with spectral index of 3.6 from hot dust emission. The C¹⁷O maps show more extended gas with the most massive molecular clump towards W3(H₂O) and none towards W3(OH) itself. We derive a peak column density of $5 \times 10^{23} \text{ cm}^{-2}$ at the H₂O maser position. Dust and line fluxes are both consistent with a mass of $10 M_{\odot}$. Peaks of molecular line channel maps of C¹⁷O, CH₃OH, and C₂H₅CN show an E-W orientation similar to the cluster of H₂O masers.

New Preprints

- 429.** Detection of CO⁺ toward the reflection nebula NGC 7023
A. Fuente, J. Martin-Pintado
1997, *ApJ Letters*
- 430.** Starburst in barred spiral galaxies – II. Molecular and optical study of three Wolf-Rayet galaxies
T. Contini, H. Wozniak, S. Considère, E. Davoust
1997, *Astronomy and Astrophysics*
- 431.** The Paschen decrement as a density indicator in MWC349*
C. Thum, A. Greve
1997, *Astronomy and Astrophysics*
- 432.** Cassegrain/Gregory-type null-correctors for surface measurements of radio telescope reflectors
A. Greve
1997, *Applied Optics*
- 433.** Structure and kinematics of a protostar: mm-interferometry of L1157
F. Gueth, S. Guilloteau, A. Dutrey, R. Bachiller
1997, *Astronomy and Astrophysics*
- 434.** Molecular gas in the spectacular ring galaxy NGC 1144
Yu Gao, P.M. Solomon, D. Downes, S.J.E. Radford
1997, *Astrophys. Journal*
- 435.** The gravitational lensing nature of the Cloverleaf unveiled in CO (7–6) line emission
D. Alloin, S. Guilloteau, R. Barvainis, R. Antonucci, L. Tacconi
1997, *Astronomy and Astrophysics*
- 436.** Multiple CO transitions, CI, and HCN from the Cloverleaf quasar
R. Barvainis, P. Maloney, R. Antonucci, D. Alloin
1997, *Astrophys. Journal*
- 437.** The chemical evolution of planetary nebulae
R. Bachiller, T. Forveille, P.J. Huggins, P. Cox
1997, *Astronomy and Astrophysics*
- 438.** Molecular gas in the barred spiral M100 – I. The IRAM 30m map
M.J. Sempere, S. Garcia-Burillo
1997, *Astronomy and Astrophysics*
- 439.** Millimeter interferometry towards the ultra-compact HII region W3(OH)
F. Wyrowski, P. Hofner, P. Schilke, C.M. Walmsley, D.J. Wilner, J.E. Wink
1997, *Astronomy and Astrophysics*
- 440.** A wideband variable waveguide coupler for millimeter applications
K.-F. Schuster, M. Carter, D. Charrière, J. Lamb, F. Mattiocco
1997, *IEEE Microwave & Guided Wave Letters*

Programmes Scheduled on the 30-m Telescope in 1996

JUNE 4 - JUNE 18

Ident.	Title	Freq. (GHz)	Authors
1.96	Time-delay measurements in the gravitational lens PDS1830-211	94	Combes, Wiklind, Kramer
121.96	High density molecular clumps in the ionized cavity surrounding the galactic center	86, 130, 217, 244	Martin-Pintado, de Vicente, Fuente, Planesas
126.96	Is SiO emission in external galaxies associated to star formation ?	86, 130, 217	Martin-Pintado, Garcia-Burillo, de Vicente, Fuente
127.96	Does dense material confine HII regions ?	108, 113, 145, 226	Martin-Pintado, Gaume, Rodriguez, de Pree, Fey
46.96	On the Fly mapping of the Cepheus-B molecular cloud	109, 115, 219, 230	Ungerechts, Guélin, Kramer, Sievers, Wild, Cernicharo
44.96	IRAS 20468 : Confirming the collapse of a low-mass protostar	98, 147, 219, 241	Wiesemeyer, Guesten
57.96	A search for redshifted [NII] 205 micron emission from BR 1202-0725	256	Van der Werf, Yun
92.96	A thorough radio spectroscopic investigation of comet Hale-Bopp	88, 96, 115, 241	Crovisier, Bockelée-Morvan, Colom, Lellouch, Rauer, Despois, Moreno
14.96	CO(1-0) along the bar and CO(1-0) (2-1) isotopes in NGC 1530	114, 109, 108	Reynaud, Downes
10.96	Determination of the CO/H2 conversion in spiral galaxies	112, 114	Kruegel, Chini

JUNE 18 - JULY 2

Ident.	Title	Freq. (GHz)	Authors
63.96	The magnetic field in the MWC349 disk (III)	86	Thum, Morris
94.96	A complete CO survey of the irregular galaxy IC 10	115,230	Wild, Kramer, Paubert, Sievers, Masset, Ungerechts, Greve
5.96	CO observations of 2 red carbon stars in the galactic halo	115, 230	Groenewegen, Oudmaijer
69.96	Does ethanol only form by evaporation from grains ?	91, 142, 234	de Vicente, Martin-Pintado
112.96	Molecular flows in the merging system of galaxies Arp299	114	Casoli, Angonin, Willaime, Gerin
113.96	CO mapping in isolated spiral galaxies	113, 115	Casoli, Sauty, Gerin

JULY 2 - JULY 7

Ident.	Title	Freq. (GHz)	Authors
122.96	The dust to gas ratio in the darkest regions of cold clouds	115, 230	Cernicharo, Cox, Zylka
123.96	Are optical jets associated to high velocity gas in molecular outflows ?	115, 230	Cernicharo, Neri, Reipurth
56.96	A conspicuous optical jet in the trifold nebula	97, 148, 230, 244	Cernicharo, Gonzalez, Cox, Lefloch, Garcia-Lopez
81.96	Probing the infrared field of the interior of hot cores	86, 87, 109, 110	Wyrowski, Schilke, Walmsley
820.96	Detection of C7H		Guelin, Cernicharo
114.96	Optical thickness of CO in Wolf-Rayet galaxies	113, 114, 226, 229	Davoust, Bridges, Wozniak, Contini, Considere
28.96	Search for S2O and OCS in IO s atmosphere	222, 219, 218	Lellouch, Belton, Strobel, Paubert
76.96	Chemical signatures of the dissipative structures of mol. clouds turbulence	89, 174, 262, 267	Falgarone, Joulain, Puget, Panis, Pineau des Forets

JULY 16 - JULY 30

Ident.	Title	Freq. (GHz)	Authors
59.96	A systematic study of the environment of Herbig Ae-Be stars	98, 110, 148, 220, 230	Fuente, Martin-Pintado, Bachiller, Palla
48.96	Molecular study of the energetic and young molecular outflow Cep E	86, 89, 96, 98, 115	Lefloch, Eisloffel, Lazareff
82.96	Bars and rings in an active early-type galaxy : NGC 4457	115, 229	Garcia-Burillo, Sempere
1.96	Time-delay measurements in the gravitational lens PDS1830-211	94	Combes, Wiklind, Kramer
84.96	The HCS+ to CS abundance ratio in cirrus clouds	85, 170, 213, 147, 244	Grossmann, Heithausen
28.96	Search for S2O and OCS in IO s atmosphere	222, 219, 218	Lellouch, Belton, Strobel, Paubert
94.96	A complete CO survey of the irregular galaxy IC 10	115, 230	Wild, Kramer, Paubert, Sievers, Masset, Ungerechts, Greve
818.96	NGC 5907		Wielebinski, Dumke

JULY 30 - AUGUST 13

Ident.	Title	Freq. (GHz)	Authors
67.96	On the Fly observations of the Trifold nebula	109, 110, 115, 230	Cernicharo, Ungerechts, Lefloch, Cox, Bachiller
36.96	Search for molecular tori around AGN		Combes, Wiklind, Drinkwater
60.96	Search for high redshift molecular absorption line systems	89, 98, 104, 147, 157	Combes, Wiklind
29.96	Molecular clouds in the dwarf elliptical galaxy NGC 205	115, 110, 230, 220	Lo, Young, lequeux
43.96	CO survey for galaxies at 0.1 z 0.5	90, 102, 160, 247	Chen, Lo

AUGUST 13 - AUGUST 27

Ident.	Title	Freq. (GHz)	Authors
87.96	Stability of clumps in Lynds 1498	109, 219, 90, 235	Gensheimer, Wilson, Lemme
92.96	A thorough radio spectroscopic investigation of comet Hale-bopp	88, 96, 115, 241	Crovisier, Biver, Boekelee, Colom, Lellouch, Rauer, Despois, Moreno
8.96	Molecular gas in a distant, dust-rich radio galaxy	87, 131	Ivison, Dunlop, Archibald, Hughes
118.96	A detailed study of molecular outflows from FU Orionis stars	110, 230, 109, 220	Eisloffel, Lefloch, Malbet
1.96	Time delay measurements in the gravitational lens PDS1830-211	94	Combes, Wiklind, Kramer
102.96	CO and ^{13}CO emission in the direction of extragalactic continuum sources	110, 115	Liszt, Lucas
11.96	A search for interstellar van der Waal complexes	86, 90, 91, 131	Havenith, Mauersberger, Wilson
7.96	CO mapping of the magellanic irregular galaxy NGC4449	115, 230	Klein, Henkel
22.96	A map of the source 3' North of Orion KL and selected positions in OMC1 in C170	112, 224, 109, 218	Wilson, Gensheimer, Dickel, Mehriger

AUGUST 27 - SEPTEMBER 10

Ident.	Title	Freq. (GHz)	Authors
11.96	A search for interstellar van der Waal complexes	86, 90, 91, 131	Havenith, Mauersberger, Wilson
1.96	Time-delay measurements in the gravitational lens PDS1830-211	94	Combes, Wiklind, Kramer
128.96	Search for the CO-H ₂ dime in the interstellar medium	91, 109	Allen, McKellar, Lequeux, Loinard
77.96	OTF mapping of M31 in CO	115, 230	Neininger, Guelin, Wielebinski, Hoernes, Berkhuijsen, Beck, Garcia-Burillo
22.96	A map of the sources 3' North of Orion KL and selected positions in OMC1 in C170	112, 224, 109, 218	Wilson, Gensheimer, Dickel, Mehriger
24.96	A new molecular core in Sgr B2	112, 224, 109, 218	Wilson, de vicente, Martin-Pintado, Gensheimer, Henkel
32.96	CN Zeeman observations : Magnetic fields in molecular clouds	113	Crutcher, Troland, Lazareff, Kazes
18.96	The ^{12}C ^{13}C ratio in the envi. of extreme ^{13}C rich carbon stars	230	Kahane, Forestini
135.95	Circumstellar dust without gas ?	115, 230	Jura, Kahane

SEPTEMBER 10 - SEPTEMBER 24

Ident.	Title	Freq. (GHz)	Authors
1.96	Time-delay measurements in the gravitational lens PDS1830-211	94	Combes, Wiklind, Kramer
78.96	The latest stages of the evolution of a bipolar outflow : NGC 7023	108, 110, 148, 220	Fuente, Martin-Pintado, Cernicharo, Rogers, Rodriguez-Franco
45.96	A CO map of the double-barred galaxy NGC 5850	114	Friedli, Combes, Leon
64.96	The giant molecular cloud W58	109, 110, 115	Thum, Ungerechts, Wink
107.96	Chemistry of the protoplanetary disks DM Tau and GG Tau	86, 144	Dutrey, Guilloteau
108.96	Chemistry of GM Aur protoplanetary disk	86, 96, 113, 226	Dutrey, Guilloteau

SEPTEMBER 24 - OCTOBER 8

Ident.	Title	Freq. (GHz)	Authors
1.96	Time-delay measurements in the gravitational lens PDS1830-211	94	Combes, Wiklind, Kramer
92.96	A thorough radio spectroscopic investigation of comet Hale-Bopp	88, 96, 115, 241	Crovisier, Biver, Bockelee et al.
77.96	On the Fly mapping of M31 in CO	115, 230	Neininger, Guélin, Wielebinski, Hoernes, Berkhuijsen, Berk, Garcia-Burillo
125.96	The eclipsed moon : Radiative and thermal behaviour of selected soils		Greve, Kramer, Pinet, Masson, Lellouch
107.96	Chemistry of the protoplanetary disks DM Tau and GG Tau	86, 144	Dutrey, Guilloteau
108.96	Chemistry of GM Aur protoplanetary disk	86, 96, 113, 226	Dutrey, Guilloteau
55.96	The origin of broad line wing emission in the Rosette molecular cloud	109, 115	Schneider, Stutzki, Williams
34.96	Multiline study of the extremely young protostellar core CB 17	86, 89, 145, 267	Launhardt, Henning, Schreyer, Ossenkopf
47.96	^{12}C ^{13}C isotopic ratio in PNe with 3He abundance determination	109, 110, 115, 230	Palla, Galli, Bachiller
58.96	Molecular gas in bipolar planetary nebulae	115, 230	Manchado, Guerrero, Bachiller
31.96	The CO- H_2 relation in planetary nebulae	115, 230	Huggins, Bachiller, Cox, Forveille
33.96	Prominent shock-chemistry variations in bipolar outflows : L1157	115, 230, 110, 220	Bachiller, Perez-Gutierrez, Tafalla

OCTOBER 8 - OCTOBER 22

Ident.	Title	Freq. (GHz)	Authors
VLBI obs.			Greve
38.96	CS towards the star forming region at the S155-Cepheus B interface	97, 146, 244	Olmi, Felli
46.96	On the Fly mapping of the Cepheus-B molecular cloud	109, 115, 219, 230	Ungerechts, Guélin, Kramer, Sievers, Wild, Cernicharo
6.96	CO search in HII galaxies	114, 228, 229	Klein, Brinks, Mebold, Heithausen, Taylor

OCTOBER 22 - NOVEMBER 5

Ident.	Title	Freq. (GHz)	Authors
49.96	Pulsar detection at short mm wavelength	90	Morris, Thum, Kramer, Wielebinski
819.96	Campaign to determine the simultaneous radio-mm-submm spectrum of SgrA*		Zylka, Falcke
6.96	CO search in HII galaxies	114, 228, 229	Klein, Brinks, Mebold, Heithausen, Taylor
96.96	Oxygen-rich chemistry in NGC 7027: An ISO/LWS follow-up study	110	Cox, Barlow, Clegg, Baluteau, Gry, Caux
1.96	Time-delay measurements in the gravitational lens PDS1830-211	94	Combes, Wiklind, Kramer
124.96	On the origin of high velocity SiO maser emission from late-type stars	86, 129, 230	Baudry, Alcolea, Cernicharo, Herpin
116.96	Compact flat-spectrum radio cores in nearby galaxies	86, 138	Reuter, Lesch
65.96	CO observations of the $z = 1.93$ hyperluminous IRAS gal. TXFSO321+009	157, 236	Roettgering, van Breugel, de Breuck, Dey
17.96	Probing shocks and fast wind in the bipolar flow of V Hya.	88, 147, 244, 130	Kahane, M-Morris, Barnbaum
120.96	CN and HCO+ in the butterfly NGC2346	226, 89	Bremer, Neri
40.96	OTF CS survey of the gal. center cloud: interaction dense molecular gas/nonthermal filament	97, 146, 244	Kramer, Staguhn, Ungerechts, Lefloch, Sievers, Paubert
106.96	Search for the HC ₄ NC isomer of HC ₅ N in molecular and circumstellar clouds	84, 86	Guelin, Cernicharo, Thaddeus, Gottlieb
90.96	The zero spacing flux in the ¹² CO(1-0) Bure observations towards the center of NGC 891	115	Garcia-Burillo, Guelin
47.96	¹² C ¹³ C isotopic ratio in PNe with 3He abundance determination	109, 110, 115, 230	Palla, Galli, Bachiller

NOVEMBER 5 - NOVEMBER 19

Ident.	Title	Freq. (GHz)	Authors
47.96	^{12}C ^{13}C isotopic ratio in PNe with ^3He abundance determination	109, 110, 115, 230	Palla, Galli, Bachiller
120.96	CN and HCO+ in the butterfly NGC2346	226, 89	Bremer, Neri
73.96	^{13}CO emission from molecular complexes in M33	110	Viallefond, Guelin, Cox
37.96	Towards the IMF : Protostellar condensations in cluster forming cores	109, 96	Blitz, Williams
1.96	Time-delay measurements in the gravitational lens PDS1830-211	94	Combes, Wiklind, Kramer
92.96	A thorough radio spectroscopic investigation of comet Hale-Bopp	88, 96, 116, 241	Crovisier, Biver, Bockelee-Morvan, Colom, Lellouch, Rauer, Despois
141.96	Time delay measurements in the gravitational lens PKS1830-211		Combes, Wiklind, Kramer
79.96	C^{17}O observations of the molecular surroundings of W3(OH)	109, 112	Wyrowski, Hofner, Walmsley, Wink
194.96	C^{34}S observations of the clumps in Lynds 1498	96, 144, 244	Gensheimer, Wilson, Lemme
196.96	H_2CO and SiO observations of NGC 2024	86	Wilson, Wink, Gensheimer, Mauersberger, Walmsley
204.96	Physics and chemistry of a newly discovered molecular core in the SgrB2 region	91, 147, 220, 112	Wilson, Gensheimer, Martin-Pintado, de Vicente
248.96	M81 : a standard galaxy ?	Bolometer	Brouillet, Baudry, Combes, Kaufman, Bash

NOVEMBER 19 - DECEMBER 3

Ident.	Title	Freq. (GHz)	Authors
216.96	CO(1-0) along the bar and CO(1-0 and (2-1)isotopes in the central concentration of NGC 1530	114, 109, 108	Reynaud, Downes
173.96	Further 30m telescope observations of OH231, 8+4.2	110, 115, 147, 230	Bujarrabal, Alcolea, Contreras
141.96	Time delay measurements in the gravitational lens PKS1830-211		Combes, Wiklind, Kramer
260.96	Excitation of a molecular cloud by the supernova remnant 3C391	98, 147, 244, 230	Reach, Rho, Wilner
158.96	Carbon, nitrogen and oxygen isotopes in the molecular envelopes of evolved stars	86, 92, 138, 224	kahane, Forestini, Guelin, Cernicharo
190.96	Isotopic ratios of heavy elements in evolved stars	89, 145, 163, 176	Kahane, Forestini, Guelin, Cernicharo
140.96	Probing shocks in the Wolf-Rayet nebula NGC 2359	110, 115, 147, 230	Kahane, St Louis, Doyon
238.96	^{13}CO and C^{18}O in the nuclear region of IC 342	109, 110, 230	Viallefond, Van Trung, Rieu

DECEMBER 3 - DECEMBER 17

Ident.	Title	Freq. (GHz)	Authors
238.96	^{13}CO and C^{18}O in the nuclear region of IC 342	109, 110, 230	Viallefond, Van Trung, Rieu
190.96	Isotopic ratios of heavy elements in evolved stars	89, 145, 163, 176	Kahane, Forestini, Guelin, Cernicharo
141.96	Time delay measurements in the gravitational lens PKS1830-211		
144.96	Observations of the hot gas in clusters of galaxies with Diabolo through the Synyaev-Zel'Dovich effect	Bolometer	Desert, Bernard, Delabrouille et al.
145.96	Anisotropy measurements of the cosmic microwave background with Diabolo at the arcmin scale	Bolometer	Desert, Bernard, Delabrouille et al.
151.96	Dust properties of cold cores in molecular clouds : A diabolo study	Bolometer	Giard, Gaertner, Ristorcelli, Serra, Andre
162.96	Coordinated centimetric and millimetric observations of radio emitting X-ray binaries	Bolometer	Paredes, Mirabel, Marti, Peracaula
152.96	Emission mechanisms in quasars	Bolometer	Chini, Kreysa, Meisenheimer, Klaas
154.96	Dust at high z ?	Bolometer	Chini, Kruegel, Sievers
254.96	Systematic study of 1.25mm emission of radioquiet QSOs with z 4	Bolometer	Omont, McMahon, Cox, Bergeron, Kreysa

DECEMBER 17 - DECEMBER 31

Ident.	Title	Freq. (GHz)	Authors
162.96	Coordinated centimetric and millimetric observations of radio emitting X-ray binaries	Bolometer	Paredes, Mirabel, Marti, Peracaula
254.96	Systematic study of 1.25mm emission of radioquiet QSOs with z 4	Bolometer	Omont, McMahon, Cox, Bergeron, Kreysa
218.96	The physics of the NGC 1333/IRAS2 eastern and western shocks	86, 91, 109, 219	Castets, Lefloch, Langer
132.96	Monitoring of Jupiter after the comet SL9 collision	88, 115, 146	Marten, Moreno, Paubert
134.96	Molecular gas in the chemically young galaxy Mrk 109	111, 115	Frayser, Sauvage, Thuan, Seaquist
141.96	Time delay measurements in the gravitational lens PKS1830-211		Combes, Wiklind, Kramer
244.96	Chemical bistability in dark clouds : the diagnostic of deuterium fractionation	86, 90, 152, 262	Gerin, Falgarone, Roueff, Le Bourlot, Pineau des Forets
45.96	A CO map of the double-barred galaxy NGC 5850	114	Friedli, Combes, Leon
823.96	Search for the primordial molecule LiH in a dense molecular cloud		Combes, Wiklind
240.96	A thorough radio spectroscopic investigation of comet Hale Bopp	88, 96, 115, 241	Crovisier, Biver, Bockelee, Lellouch et al.
206.96	Molecular gas in faint blue galaxies at intermediate redshifts	153, 158, 138, 151	Wilson-C, Combes

The IRAM Newsletter is edited by Robert LUCAS at IRAM-Grenoble (e-mail address: lucas@iram.fr).

The IRAM Newsletter is available in electronic form:

- by using the World Wide Web: from the IRAM home page (<http://iram.fr/>), click on item "Newsletter" and follow the links...

- by means of an anonymous ftp account, opened at IRAM for Internet users. To access those files, please connect through ftp to [iram.fr](ftp://iram.fr) (or 193.48.252.22) and read the README file. Several subdirectories are available:

Directory	Contents
/dist/newsletter	Recent issues of this Newsletter (one subdirectory per issue)
e.g. /dist/newsletter/jul95	jul95.ps is the Postscript file for the July 1995 issue.
/dist/doc	Documentation on IRAM telescopes and software
/dist/proposal	Proposal forms and Latex files to aid proposal preparation
/dist/soft	distribution files for reduction software

- by means of an electronic mail file server installed at IRAM (on the Alpha machine IRAM04). This file server is a file distribution service that uses electronic mail facilities to deliver files. To communicate with it you should send a message to the electronic address:

newsserv@iram.grenet.fr

For instance, to obtain a copy of the May 1992 issue, just send the one line message:

SENDME MAY92.PS

to the above electronic address. You will receive later a mail message containing the IRAM Newsletter in Postscript code. Please discard all the e-mail header information with a text editor, and send the file to a Postscript laser printer.

More information may be obtained by sending the one line message:

HELP

Note that this file server also contains Postscript files of the proposal forms and of Plateau de Bure documentation.

We also compile a list of e-mail addresses of IRAM users (e.g., in order to send warning messages when the Newsletter is available, but also to provide fast information, if needed). If you feel your address should be on this list, please send the one line message:

SUBSCRIBE NEWSSERV *your name*

to the following e-mail address:

newsserv@iram.grenet.fr

Both addresses are valid on Internet, EARN-Bitnet and EAN Please keep R. Lucas informed of any problem you may encounter.

IRAM Addresses:

	Address:	Telephone:	Fax:
Grenoble	Institut de Radioastronomie Millimétrique, 300 rue de la Piscine, Domaine Universitaire, 38406 St Martin d'Hères Cedex, France		
	from abroad:	33 476 82 49 00	33 476 51 59 38
	from France:	0 476 82 49 00	0 476 51 59 38
Plateau de Bure	Institut de Radioastronomie Millimétrique, Observatoire du Plateau de Bure, 05250 St Etienne en Dévoluy, France		
	from abroad:	33 492 52 53 60	33 492 52 53 61
	from France:	0 492 52 53 60	0 492 52 53 61
Granada	Instituto de Radioastronomía Milimétrica, Avenida Divina Pastora 7, Núcleo Central, 18012 Granada, España	(34) 58 22 88 99	(34) 58 22 23 63
Pico Veleta	Instituto de Radioastronomía Milimétrica, Estación Radioastronómica IRAM-IGN del Pico Veleta, Sierra Nevada, 18012 Granada, España		
	starting April 15:	(34) 58 48 20 02	(34) 58 48 11 48

E-Mail Addresses:

- IRAM-Grenoble: username@iram.fr

- IRAM-Granada: username@iram.es

The **username** is generally the last name of the person to be contacted.